




ISSN : 2347-2251
**Indo-American Journal of
Pharma and Bio Sciences**



www.iajpb.com

iajpb.editor@gmail.com
editor@iajpb.com



PRELIMINARY PHYTOCHEMICAL AND ANTIMICROBIAL SCREENING OF SEED AND COAT OF CITRUS SINENSIS

¹Mrs. V. Padmaja,²Dr. C. Vijitha,³Mr. A. Ramesh

¹Professor,^{2,3}Associate Professor

^{1,2}Department of Pharmacognosy

³Department of Pharm.Chemistry

Vaagdevi Institute of Pharmaceutical Sciences, Bollikunta, Warangal. Telangana.

ABSTRACT:

One of the most widely consumed citrus species is *Citrus sinensis*. Orange peels and seeds are waste products that come from the fruit and might have valuable phytonutrients that are relevant to biology.

Sweet orange peels, both fresh and dry, were macerated. Using accepted techniques, the total alkaloid, flavonoid, and tannin content was calculated. Using the disc diffusion method, antimicrobial activities were observed against five bacterial strains (*Staphylococcus aureus*, *Enterococcus faecalis*, *Pseudomonas aeruginosa*, *Escherichia coli*, and *Salmonella typhimurium*) and three fungal strains (*Candida albicans*, *Aspergillus niger*, and *Penicillium notatum*). The fresh peel extract had a higher total alkaloid, flavonoid, and tannin content than the dry peel extract. Comparing fresh and dry peel extracts, antimicrobial activities showed that the former had superior antibacterial activity against every bacterial strain and one fungus strain under investigation. On the other hand, the dried peel extract was more effective than the fresh peel extract in inhibiting the growth of *Aspergillus niger* and *Penicillium notatum*. This study looked at the antibacterial properties and phenolic content of peel extracts from fresh and dried *Citrus sinensis*. The study's findings

indicate that, in comparison to the dried peel extract, the fresh *Citrus sinensis* peel extract has more phenolic content and stronger antibacterial activity against the microbial strains under investigation. The results of this study imply that phytochemicals may be lost if plant components are dried before being extracted for phytonutrients.

KEYWORDS: *Citrus sinensis*, Orange, Antimicrobial, Phytochemicals, Seeds, Peels.

1. INTRODUCTION:

Citrus sinensis (L.) Osbeck (sweet orange) is the world's most widely grown and commercialized citrus specie. The fruit of *C. sinensis* is mostly recognized for its vitamin C content and is also an important source of other phytochemicals such as phenolics and carotenoids which are reputed to have health benefits. The sweet orange fruit is usually eaten whole or processed into juice after the peeling of the external rind (flavedo). This peeling process leads to the generation of substantial wastes. Sweet orange (*Citrus sinensis* (L.)) is the world's most commonly cultivated fruit tree. It belongs to the Rutaceae family which comprises mandarins, limes, lemons, grapefruits, sour and sweet orange. Citrus fruits are of



immense economic value; occupying the top position in fruit production. Orange trees are widely cultivated in tropical and subtropical climates for the sweet fruit, which is peeled or cut (to avoid the bitter rind) and eaten whole, or processed to extract orange juice. Citrus fruits are known to contain substantial quantities of vitamin C, a potent water-soluble vitamin essential for healthy living. The term phytochemical is often used to describe a diverse range of biologically active compounds found in plants. Phytochemicals provide plants with colour, flavour and natural protection against pests. Phytochemicals are not essentially required for the sustenance of life but confer extra health benefits against pathogens. Phytochemical screening avails us with the opportunity to see at a glance the various phytochemicals present in a plant material. This may give a hint as to the possible range of bioactivities the plant product may possess. It also serves as a preliminary step in research protocol aimed at the isolation, purification and utilization of compounds inherent in the plant material for medical, pharmaceutical or agro-industrial use. The orange fruit is of unique economic importance as all portions contain potentials for diverse industrial usability. Thus far, researches into citrus wastes have been concentrated on the peels (flavedo); with little interest on the albedo and seeds. This study aims to draw attention to the possible waste-to-wealth utilization of orange fruit by-products with the added advantage of providing novel sources of nutraceuticals, phytochemicals, pharmaceuticals and reducing environmental insults.

2. PLANT PROFILE:

The *Citrus sinensis* popularly known as sweet orange seed in Igbo of Nigeria is of the Rutaceae family. The seed is best sown

in a greenhouse as soon as it is ripe after thoroughly rinsing it, sow stored seed in March in a greenhouse, germination usually takes place within 2-3 weeks at 130C. seedlings are liable to damp off so they must be watered with care and kept well ventilated. *Citrus sinensis*. Contains a wide range of active ingredients and research is still underway in finding uses for them. They are rich in vitamin C, flavonoids, acids and volatile oils. They also contain coumarins such as bergapten which sensitizes the skin to sunlight. Bergapten is sometimes added to tanning preparations since it promotes pigmentation in the skin, though it can cause dermatitis or allergy responses in some people. Some of the plants more recent applications are as sources of anti-oxidants and chemical exfoliants in specified cosmetics. The fruit is an appetizer and blood purifier, it is used to allay thirst in people with fever and also treat catarrh. The fruit juice is useful in treatment of bilious infections and bilious diarrhea. The fruit rind is caminative and tonic cure for acne. The dried peel is used in the treatment of anorexia, cold cough etc.

Origin And Description :

The orange is unknown in the wild state; its assumed to have originated in Southern China. Northeastern India and perhaps Southeastern Asia (formally Indochina). It was carried to the mediterenian area possibly by Italian traders after 1450 of by Portuguese navigators around 1500. Up to that era citrus fruits were valued by Europeans mainly for medicinal purposes, but orange was quickly adopted as a luscious fruit and wealthy persons grow it in private conservations, called orangeries. By 1646, it had been much publicized and was well known. The orange has become the most commonly grown fruit in the world. It is an important crop in the far east,



the union of South Africa, Australia, throughout the Mediteranian area and sub tropical areas of South America and the Caribbean. The United States leads in the world production, with Florida, alone, having an annual yield of more than 200 million boxes, except when freezes occur which may reduce the crop by 20 or even 40%.

Morphology Aspect Of The Citrus Sinensis :

The orange tree, reaching 25 ft (7.5m) or with great ages up to 50ft (15m) has a rounded crown of slender branches. The twigs are twisted and angled when young and may bear slender semi- flexible, bluntish spines in the leaf axils. There may be faint or conspicuous wings on the petioles of aromatic evergreen, alternate elliptic to ovate, sometimes faintly toothed “leaves” – technically solitary leaflets of compound leaves. These are 2 1/2 to 6m (6.5-15cm) long 1 to 3 3/4 in (2.5 – 9.5cm) wide. Brone singly or in clusters of 2 to 6, the sweetly fragrant white flowers, about 2 in (5cm) wide, have a saucershaped, 5 pointed calyx and 5 oblong, white petals, and 20 to 25 stamens with conspicuous yellow anthers. The fruit is subglobose oblate or some what oval, 2 1/2 to 3 3/4 in (6.5-9.5cm) wide. Dotted with minute glands containing an essential oil, the outer ring (epicarp) is orange or yellow when ripe, the inner ring (mesocarp) is white spongy and nonaromatic, the pulp (endocarp) yellow, orange or more less red, the sweet orange differs physically from sour orange in having a solid center.

Phytoconstituents: 1.5 % essential oil is present in the orange fruit. D-limonene (90%), citral, sinesal, n-nonanal, n-decanal, ndodecanal, geranyl acetate, anthranil acid,

citronellal, linalyl acetate, methyl ester are present.

SR.	PHYTOCONSTITUENTS	PLANT PART
1	Flavone glycosides; Neohesperidin, Naringin, Hesperidin, Narirutin Triterpene; Limonene, Citrol Pigment; Anthocyanin, Beta-cryptoxanthin, Cryptoxanthin, Zeaxanthin and Rutin, Eriocitrin, Homocysteine Polymethoxylted flavones; Tangeritin and Nobiletin Flavonoids; Citricidone, Citabrsine and Noradrenaline	Fruit peel
2	Terpanoids, Linalool, β elemene	Leaves
3	Triterpenes, Limonene	Flowers
4	Vitamins; B1, B2, B3, B5, B6, and Vitamin C Minerals;	Fruits

3. MATERIAL METHOD:

The peel part of citrus sinensis was collected and dried under shade. This dried material was mechanically powdered, sheaved and stored at a dry place. This powdered material was used for further phytochemical, antimicrobial and antioxidant analysis.

Collection of plant material:

The fresh peels of citrus sinensis were collected from the local market in Bhopal. The peels were washed thoroughly with tap water and dried under shade for 7-10 days grounded into fine powder and sieved.

Preparation of powder: The peel part of citrus sinensis was collected and dried under shade. These dried materials were mechanically powdered, sheaved using 80 meshes and stored in an airtight container. These powder materials were used for further phytochemical analysis.

Extraction of the plant material : The peel of the plant were properly washed in tap water and rinsed in distilled water. The rinsed Peel part were hot air-dried for 7-10 days at room temperature. The dried peel part of each plant were pulverized using pestle mortar to obtain a powdered form which was stored in airtight glass containers at 4°C until used. 50 g of powdered sample was soaked in distilled water and methanol (200 mL and 100 mL)



separately for 12 hrs at room temperature. The extracts were then filtered with whatsmann filter paper no 4 and concentrated to a final volume of 50 mL and subjected to phytochemical analysis.

Phytochemical Screening :

The phytochemical screening involves the simple chemical test to detect the presence of secondary metabolites.. The phytochemical test include: tests for saponins, tannins, flavonoids, alkaloids, cardiac glycosides and phenol.

Test for Alkaloids : 12 ml of chloroform was stirred with peel extracts separately with a few drops of dilute hydrochloric acid and filtered. The filtrates were divided into four equal portions and tested with various alkaloidal reagents namely Mayer's reagent (cream precipitate), Dragendorff's reagent (orange brown precipitate), Wagner reagent (reddish brown precipitate), and Picric acid (white precipitate).

Test for Saponins :

Frothing test: 0.5 g of the extract was weighed and dissolved in 10 ml of sterile distilled water. The test tube was stopped and shaken vigorously for 30 seconds. It was then allowed to stand for half an hour.

Foaming test: 0.5 g of the extract was weighed and stirred in 20 ml of sterile distilled water and boiled in a water bath for 5 minutes and filtered. 10 ml of the filtrate was mixed with 5 ml of distilled water and shaken vigorously for a stable persistent froth. The frothing was mixed with 3 drops of olive oil and shaken vigorously, then observed for the formation of emulsion.

Sodium bicarbonate test: 0.5 g of the extract was weighed and dissolved in 10 ml of sterile distilled water. 3 drops of sodium bicarbonate solution were added and

shaken vigorously for two minutes, then observed for the formation of emulsion.

Test for Flavonoids :

Ammonia test: Filter paper strips were dipped in the ethanolic solutions of the extract and ammoniated. The filter paper changed its colour to yellow which indicates the presence of flavonoids.

Shinoda's test: To 1 ml of the each extract, a piece of metallic magnesium was added followed by addition of 2 drops of concentrated hydrochloric acid. A red colour confirmed the presence of flavonoids in the extract.

Sodium hydroxide test: Aqueous sodium hydroxide was added to 1 ml of the extract solution, dilute sulphuric acid was added slowly. A yellow colour which turned cream on addition of dilute sulphuric acid confirmed the presence of flavonoids.

Test for Tannins : Ferric chloride test : About 0.5 g of the extract was stirred with 2 ml of distilled water and filtered. The filtrate was treated with five percent (5 %) ferric chloride reagent. A blue-green precipitation was taken as evidence for the presence of tannins.

Bromine water test: About 0.5 g of the extract solution was dissolved in distilled water in a test-tube. 5 drops of bromine water was added gently. Decolouration of the bromine water confirmed the presence of tannins.

Test for Cardiac Glycosides : Three grammes (3 g) of the extract were hydrolysed with dilute hydrochloric acid for 2 hours in water bath and allowed to cool well on ice for 2 minutes and filtered. The filtrate was subjected to different test for cardiac glycosides: Liebermann-



burchard's, Keller-killani, and Salkowski tests.

Keller-Killani Test: One millilitre (1 ml) of glacial acetic acid containing traces of FeCl_3 and one millilitre (1 ml) of concentrated H_2SO_4 was added to the extract carefully. The formation of brown ring at the interface indicated the presence of cardiac glycosides.

Salkowski test: To another portion of the solution, one millilitre (1 ml) of sulphuric acid was added carefully to form lower layer. A red brown colour at the interface indicated the presence of cardiac glycosides.

Liebermann's test: To another portion of the solution, one millilitre (1 ml) of hydrochloric acid was added carefully to form lower layer. A green colour with a ring formation on top while blue-green colour indicated the cardiac glycosides.

Test for phenols : One milliliter of the extract was added to 1ml of 10% FeCl_3 and mixed together. The presence of blue precipitate confirms the presence of Phenols.

Screening for Antimicrobial Activity : The hot and cold aqueous and ethanol extracts of the citrus sinensis were used for the antimicrobial screening using the agar well diffusion method. The media was punched with 7 mm diameter wells and were filled with various concentrations of the extracts 5 mg/ml, 10 mg/ml, 15 mg/mL, 20 mg/ mL and 25mg/mL. The plates were then incubated at 37°C for 24 hours. After incubation, zone of growth inhibition for each extract was measured in millimeters by using a scale.

4. RESULT AND DISCUSSION:

Observation Table of Phytochemical Screening

S.NO	TEST	RESULT	INFERENCE
1.	Alkaloids Test		
	Dragendroff Test	Present	Yellow / brown ppt
	Mayers Test	Present	Cream colour obtain
	Wagners Test	Present	Redish Brown ppt
2.	Saponins Test		
	Foaming Test	Present	Formation of emulsion
	Sodium Bicarbonate Test	Absent	Formation of emulsion
3.	Flavonoids Test		
	Ammonia Test	Absent	Yellow colour obtain
	Shinoda Test	Present	Red colour obtain
	Sodium Hydroxide Test	Present	Yellow colour obtain
4.	Tannin Test		
	Ferric Chloride Test	Present	Blue green colour obtain
	Bromine Water Test	Absent	Decolouration of Bromine Water
5.	Cardiac Glycosides Test		
	Keller Kiliani Test	Present	Brown Ring obtain
	Liebermann's Test	Present	Blue green colour obtain
	Salkowshi Test	Absent	Red brown colour obtain
6.	Phenol Test	Present	Blue colour obtain

The results show that both the FPE and DPE of *C. sinensis* possess varying degrees of antimicrobial activities against the test bacterial and fungal strains (Tables 1 and 2). The FPE produced the widest zone of inhibition (ZOI) of 20 mm against *E. faecalis*. This was followed by *S. aureus* and *P. aeruginosa* with 14 mm ZOI and *E. coli* with 13 mm ZOI (Table 1). The FPE produced a 6 mm ZOI for *S. typhimurium*., the lowest observed for the bacterial strains studied. The DPE produced generally smaller zones of inhibition against the bacterial strains with a 12 mm zone of inhibition observed for *E. faecalis* and 10 mm for *S. typhimurium*. The DPE produced 4 mm, 6 mm and 8 mm zones of inhibition, respectively against *S. aureus*, *P. aeruginosa* and *E. coli*.

Table 1: Antibacterial activities of *Citrus sinensis* peel extracts (200 $\mu\text{g/mL}$) against some bacterial strains tested by disc diffusion assay



	Zones of inhibition (mm)				
	Gram positive		Gram negative		
	<i>Staphylococcus aureus</i>	<i>Enterococcus faecalis</i>	<i>Pseudomonas aeruginosa</i>	<i>Escherichia coli</i>	<i>Salmonella typhimurium</i>
FPE	14	20	14	13	6
DPE	4	12	6	8	10

Values are mean of 3 biological replicates to the nearest mm.

Table 2: Antifungal activities of *Citrus sinensis* peel extracts (200 µg/mL) against some bacterial strains tested by disc diffusion assay

	Zones of inhibition (mm)		
	<i>Candida albicans</i>	<i>Aspergillus niger</i>	<i>Penicillium notatum</i>
FPE	18	2	2
DPE	2	4	10

Values are mean of 3 biological replicates to the nearest mm

Table 2 shows that the FPE was most effective against *C. albicans*, producing an 18 mm ZOI while the DPE was most effective against *P. notatum*. with an observed ZOI of 10 mm. Two 2 mm zones of inhibition were observed for the FPE against *A. niger* and *P. notatum*. while 2 and 4 mm respectively for *C. albicans* and *A. niger* when exposed to the DPE.

5. SUMMARY & CONCLUSION:

The foundation for future investigation into the identification of the antibacterial substances found in *C. sinensis* peels is this preliminary study. This highlights even more the waste-to-wealth potential of wastes from sweet oranges. The findings of this investigation demonstrate that, in comparison to the dried peel extract, the fresh *C. sinensis* peel extract has higher phenolic content and superior antibacterial activity against the pathogens under investigation.

Our results also imply that it might not always be a good idea to dry plant materials before extracting them because some of the active pharmacological chemicals might be lost in the process.

REFERENCES:

- Oikeh EI, Oriakhi K, Omoregie ES. Phenolic content and in vitro antioxidant activities of sweet Orange (*Citrus sinensis* L.) fruit wastes. *Archives of Basic and Applied Medicine*. 2014;2(2):119–26.
- Liu Y, Heying E, Tanumihardjo SA. History, global distribution, and nutritional importance of Citrus fruits. *Compr Rev Food Sci Food Saf*. 2012;11(6):530–45.
- Martínez-Cuenca MR, Primo-Capella A, Forner-Giner MA. Influence of rootstock on Citrus tree growth: effects on photosynthesis and carbohydrate distribution, plant size, yield, fruit quality, and dwarfing genotypes. 2016. InTech.
- Mandalari G, Bennett RN, Bisignano G, Saija A, Dugo G, et al. Characterization of flavonoids and pectin from bergamot (*Citrus bergamia* Risso) peel, a major byproduct of essential oil extraction. *Journal of Agriculture and Food Chemistry*. 2006;54:197–203.
- Hegazy AE, Ibrahim MI. Antioxidant activities of Orange Peel extracts. *World Appl Sci J*. 2012;18(5):684–8.
- Putnik P, Bursać Kovačević D, Režek Jambrak A, Barba FJ, Cravotto G, et al. Innovative "Green" and Novel Strategies for the Extraction of Bioactive Added Value Compounds from Citrus Wastes A Review. *Molecules* (Basel, Switzerland), 2017;22(5), 680.
- Oikeh EI, Oriakhi K, Omoregie ES. Proximate analysis and phytochemical



screening of *Citrus sinensis* fruit wastes. *The Bioscientist*. 2013;1(2):164–70.

8. Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety. *Front Pharmacol*. 2014;4:177.

9. Yuan H, Ma Q, Ye L, Piao G. The traditional medicine and modern medicine from natural products. *Molecules*. 2016;21(5):559.

10. Onyeagba RA, Ugbogu OC, Okeke CU, Iroakasi O. Studies on the antimicrobial effects of garlic (*Allium sativum* Linn), ginger (*Zingiber officinale* roscoe) and lime (*Citrus aurantifolia* Linn). *Afr J Biotechnol*. 2004;3(10):552–4.

11. Ayukekbong JA, Ntemgwa M, Atabe AN. The threat of antimicrobial resistance in developing countries: causes and control strategies. *Antimicrob Resist Infect Control*. 2017;6:47.